



Ankle: Weber A or avulsion fracture

— Virtual Fracture Care

You have a fracture of the ankle (weber A) or a "pull-off" fragment of one of the bones in your foot (avulsion fracture). These fractures are similar to a torn ankle ligament and are treated in the similarly.

Summary

- You have a fracture of the ankle or a small 'pull-off' fragment of one of the bones in your foot.
- This type of injury almost always heals well without any loss of function.
- A routine follow-up is not necessary for this type of injury.
- It is important that you keep your ankle elevated as much as possible during the first week, this will reduce the swelling.
- Wear the brace as soon as the swelling has sufficiently decreased. After three weeks, try wearing the brace less often, if your pain allows you to do this.
- You can find exercises for your ankle in the folder or in the app.
- If you have any questions, or are experiencing problems with your recovery, you can call the "Breuklijn" (fracture hotline) on work days between 08:00 – 10:00 and 13:00 – 15:00.

Treatment

A Weber A or avulsion fracture of the ankle is a common injury. The fracture is treated with a tubigrip (an elastick sock) and an ankle brace. You can rent crutches at the hospital.

0-1 week

- Tubigrip.
- Wear the brace as soon as the swelling has sufficiently decreased to allow you to do this (usually after 3-4 days).

1-3 weeks

- Wear the tubigrip and brace.

3-6 weeks

- Try wearing the brace less often, as soon as your pain allows this.

After 6 weeks

- You may start sports.

Instructions

Follow these instructions for a good recovery.

- You can combine wearing the ankle brace and a sturdy shoe, for example a shoe used for hiking. You can stand on the foot as long as the pain allows you to do this.
- You can gradually reduce the use of the brace after 3 to 6 weeks. You can find instructions on how to reapply the brace in the app.
- It is important to elevate your ankle. Does the brace feel more tight? Elevate the ankle, for example, by putting it on a chair. The swelling will reduce, and the pressure and pain will decrease.
- The fracture is close to the joint. Start moving the ankle as soon as possible, to prevent the ankle from getting stiff. You can find exercises in the app or in the information folder. It is important not to do the exercises if your pain does not allow this.
- For the first six weeks, you should not practice any sports. After six weeks, you may start sports if your pain allows you to do this. You can wear a brace during sports if necessary.
- Are you in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait with the use of paracetamol until the pain is too much.

Recovery

- This type of injury almost always heals well with time and use – no specific treatment is required and routine follow-up is not necessary.
- Physiotherapy is not necessary. If you are not satisfied with the function of your ankle after 6 weeks, you can contact a physiotherapist. The app contains exercises to recover better.
- Does the pain increase, or does it not improve? Please contact the "Breuklijn" (fracture hotline).

Reapply the brace

If you want to reapply the brace, you can find instructions in the video below to do so.

Video: [Aanleggen enkelbrace bij enkelbandletsel / weber A - YouTube](#)

Exercises

After a fracture of your ankle, stiffness, loss of strength and loss of coordination may occur. The following exercises can be performed to support your recovery.

When to start and what to do

You can start doing exercises after four weeks. Start with non-weight bearing exercises. This means that you perform exercises without standing on your foot. When you are able to successfully perform these exercises, you can start with weight-bearing exercises. Only exercise as pain allows.

Follow the instructions below:

- Practice at least 3 times a day and repeat each exercise 10 times
- you can perform the exercises in warm water if preferred

Non-weight bearing exercises

Non-weight bearing exercises can be divided into passive and active exercises. Start with the passive exercises and then continue with the active exercises.

- **Passive exercises:** bend, straighten and move the ankle with your hands.
- **Active exercises:** without using your hands, tighten the calf and foot muscles, bend, straighten and move the ankle in all directions.

Weight bearing exercises

Weight bearing exercises can also be divided into passive and active exercises. Again, start with the passive exercises and then continue with the active exercises.

- **Passive exercises:** sit down and do the stretching exercises using your body weight, as shown in the video below.

Video: [Enkel extensie oefening - YouTube](#)

- **Active exercises:** stand up and do the stretching exercises using your body weight, as shown in the video below.

Video: [enkel extensie oefening - YouTube](#)

Other exercises

It is important to practice standing correctly on the affected leg, as pain allows. Walking starts with being able to stand on one leg. Also try to use the ankle as normally as possible again, as pain allows. Every step you take is a step closer to a normal gait.

Questions?

You can call us.

If there are any questions after reading the information, please contact us or visit www.spaarnegasthuis.nl

Contact

Breuklijn: (023) 224 0025

Please call us on work days between 08:00 – 10:00 and 13:00 – 15:00.

Directly to the information in the Virtual Fracture Care app?

The QR code below will take you directly to the information from this folder.



Waar zijn we te vinden?

Haarlem Zuid

Boerhaavelaan 22
2035 RC Haarlem

(023) 224 0000

Haarlem Noord

Vondelweg 999
2026 BW Haarlem

www.spaarnegasthuis.nl

Hoofddorp

Spaarnepoort 1
2134 TM Hoofddorp

info@spaarnegasthuis.nl