Collarbone fracture (children)

Virtual Fracture Care

Your child has a fractured collarbone (clavicle). This fracture is treated with a sling.



Samenvatting

- Your child has a fractured collarbone.
- This type of injury almost always heals well without any loss of function.
- A routine follow-up is not necessary for this type of injury.



- You can remove the sling as soon as your child's pain allows you to do this. During the night you do not have to wear the sling. Use the sling for max. 4 weeks
- To prevent the shoulder from getting stiff, it is important that your child starts moving the shoulder and arm as soon as possible. Only use the shoulder when the pain allows this.
- You can find instructions on how to reapply the sling in the app.
- If you have any questions, or are experiencing problems with your recovery, you can call the "Breuklijn" (fracture hotline) on work days between 08:00 10:00 and 13:00 15:00.

Treatment

A fractured collarbone is a common injury in children. The fracture is treated with a sling. If your child is young ans won't wear the sling, this is not a problem. Collarbone fractures almost always heal without any problems or complications.

0-4 weeks

• Wear the sling, remove it as soon as your child's pain allows this.

4-6 weeks

• Let your child move the shoulder if the pain allows this.

After 6 weeks

• Gradually start sports.

Instructions

Follow these instructions for a good recovery.

- The sling keeps your child's arm in a comfortable position for a few days to weeks.
- You can remove the sling as soon as your child's pain allows you to do this. During the night you do not have to wear the sling. Use the sling for a maximum of 4 weeks.
- To prevent a stiff shoulder it is important that your child starts moving his/her shoulder as soon as possible. Only use the shoulder when your child's pain allows you to do this.
- Activities such as sports can be done after 6 weeks when your child's pain allows you to do this. It is no problem to wait longer with these activities.
- Is your child in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait with the use of paracetamol until the pain is too much.

Recovery

- This type of injury almost always heals well with time and use no specific treatment is required and routine follow-up is unnecessary.
- As the bone heals, a bump may appear over the fracture. This is normal, it is the result of bone healing. This does not affect the function of your child's shoulder.
- Does the pain increase, or does it not improve? Please contact the "Breuklijn" (fracture hotline).
- Physiotherapy is not necessary. If you are not satisfied with the function of your child's shoulder after 6 weeks, you can ask for a referral to a physiotherapist from your general practitioner. However, a referral is not required.

Reapply the sling

Instructions on how to reapply the sling can be found in the video below.

Video: <u>Aanleggen sling (enkele loop) - YouTube</u>

Video: Aanleggen sling dubbel loop bij sleutelbeen breuk - YouTube

Exercises

To prevent a stiff shoulder it is important that your child starts moving the shoulder as soon as possible. Only use the shoulder when the pain allows you to do this. No specific exercises are necessary.

Questions?

You can call us. If there are any questions after reading the information, please contact us or visit www.spaarnegasthuis.nl

Contact

Breuklijn: (023) 224 0025 Please call us on work days between 08:00 – 10:00 and 13:00 – 15:00.

Directly to the information in the Virtual Fracture Care app?

The QR code below will take you directly to the information from this folder.



Waar zijn we te vinden?

Haarlem Zuid Boerhaavelaan 22 2035 RC Haarlem

(023) 224 0000

Haarlem Noord Vondelweg 999 2026 BW Haarlem

www.spaarnegasthuis.nl

Hoofddorp Spaarnepoort 1 2134 TM Hoofddorp

info@spaarnegasthuis.nl